

## Production of Fermented Beverage from Soursop Fruit

Nguyen Phuoc Minh\*

Tra Vinh University, Vietnam

\*Corresponding Author E-mail: [dr.nguyenphuocminh@gmail.com](mailto:dr.nguyenphuocminh@gmail.com)

### ABSTRACT

*Soursop, Annona muricata, also called guanabana is a fruit with an acidic taste, closely related to custard apple. It is highly susceptible to spoilage, softens very rapidly during ripening, and becomes mushy and difficult to consume fresh. It is sold in local markets and is rejected at market because of external injuries or uneven shape and size. We focus on the investigation of technical factors affecting to soursop juice fermentation. Our results show that pectinase 0.03%, incubation at 40°C in 2 hours, Saccharomyces cerevisiae yeast ratio 8%, pH 4.0, fermentation temperature at 26°C, fermentation time 60 hours, concentrated juice addition 30%, syrup 65°Brix with 8% supplementation, caramel 1.5%, and pasteurization at 80°C in 10 minutes.*

**Keywords:** Soursop, fermented juice, *Saccharomyces cerevisiae*, fermentation

### INTRODUCTION

Soursop fruit contains various types of nutrients beneficial to human health such as vitamins C, B1, and B12. Soursop fruit is also rich in carbohydrates, particularly fructose. Soursop has many therapeutic properties; the juice is diuretic while the other parts have antibacterial, anticancerous, astringent, sedative, and other properties<sup>2</sup>. Soursop, like other tropical fruits, serves as a potential source of raw materials for fruit products such as juice, beverages, wine, jellies, jam puree, power fruit bars, and flakes<sup>1,3,7</sup> conducted a kinetic evaluation of the fermentation of soursop (Substrate) by *Saccharomyces cerevisiae*.

Raphael N Okigbo and Omokaro Obire (2009) conducted on the mycoflora associated with the different parts of fresh and rotten fruits of soursop (*Annona muricata* L.) and the potential of using both indigenous yeast flora and commercial yeast extract for wine production. Sunday P. Ukwo, Chidi F. Ezeama (2011) investigated the proliferation of acetic acid bacteria during various stages of soursop juice fermentation. Preliminary experiment was carried out to identify the genera and species of acetic acid bacteria associated with soursop fruits. The effect of pH, temperature and momentary aeration on the growth of acetic acid bacteria and its effect on the growth of *Saccharomyces cerevisiae* were studied. Imade, E.E. *et al.*<sup>4</sup> evaluated the changes in the microbial profile, physico-chemical and nutritional attributes during the bioconversion of soursop (*Annona muricata*) must to wine. Nwachukwu, E. and Ezeigbo, C.G. (2013) investigated the microbial quality of locally produced soursop juice and the effect of pasteurization, acidification and chemical preservative on the juice. I.E. Mbaeyi-Nwaoha, C.N. Ajumobi (2015) carried out a production and microbial evaluation of table wine from tamarind (*Tamarindus indica*) and soursop (*Annona muricata*).

The main purpose of this our research is to investigate technical conditions during soursop juice fermentation such as the yeast proliferation, pectinase, yeast ratio, pH, fermentation time and temperature etc to the quality of soursop fermented juice.

## MATERIAL AND METHOD

### Material

Fig. 1: Soursop fruit



Soursop fruits are collected in Mekong River Delta, Vietnam. They are in ripen stage without rotten or damage. *Saccharomyces cerevisiae* is supported from Pasteur Institutue, HCMC, Vietnam.

### Research method

#### Yeast proliferation

Prepare 100 ml of medium in erlen 250ml. Proliferate yeast at 25oC in 4-36 hours. By 4 hours, we take 5 ml of samples to centrifuge and count on the counter.

#### Pectinase supplementation

The soursop juice was obtained manually. Conduct the experiment with 5 samples, each sample 200ml. Investigate different pectinase ratios 0%, 0.01%, 0.02%, 0.03%, and 0.04%.

#### Sugar supplementation, pH, yeast, and time for fermentation

Soursop fruits are treated with pectinase and then filtered, adjusted pH and oBrix, added yeast. Then the juice is fermented at normal room temperature 25-27oC. During fermentation, we analyse ethanol, soluble dry matter, and total acidity. Experiment is conducted with 200 ml of each sample. Fermentation time: 24h, 36h, 48h, 60h, 72h; pH: 3.0, 3.5, 4.0, 4.5, 5.0; yeast: 4%, 6%, 8%, 10%, 12%; °Brix: 160g/l, 180g/l, 200g/l, 220g/l, 240g/l. We monitor ethanol formation, °Brix and total acidity by 12 hours.

#### Concentrated juice, syrup, caramel supplementation

Ratio of soursop concentrated juice 10%, 20%, 30%, 40%, 50%; syrup 4%, 6%, 8%, 10, 12%; caramel 0.5%, 1%, 1.5%, 2%, 2.5% are verified.

#### Pasteurization for soursop fermented juice

Soursop fermented juice bottles are pasteurized under 75°C, 80°C, 85°C, 90°C, 95°C in different durations 5, 10, 15, 20, 25 minutes.


### Statistical analysis

All data are processed by Excel

## RESULT AND DISCUSSION

### Yeast proliferation

Fig. 2: Yeast growth during the soursop fermentation



After 24 hours of fermentation, the yeast cells come to the maximum level with high activity.

**Effect of pectinase supplementation**


**Table 1. Sensory score of the soursop fermented juice**

Sample	Pectinase supplementation (%)	Sensory score of the fermented juice
1	Control (0%)	Turbid, sediment, yellow
2	0.01%	Turbid, sediment, light yellow
3	0.02%	Quite clear, sediment, light yellow
4	0.03%	Clear, low sediment, light yellow
5	0.04%	Low sediment, light yellow

With 0.03% pectinase, it's enough to hydrolyze pectin so we get the best soursop juice.

**Effect of sugar to soursop juice fermentation**


**Fig. 3: Effect of the initial sugar to fermentation**



We notice that sugar 200g/l is suitable for fermentation.

**Effect of pH for soursop fermentation**


**Fig. 4: Effect of pH to ethanol formation during soursop fermentation**



pH 4 is optimal for soursop fermentation

**Effect of yeast ratio for soursop fermentation**


**Fig. 5: Effect of yeast ratio to soursop fermentation**



We see that 5% yeast is appropriated for soursop fermentation

**Effect of time to soursop fermentation**


**Fig. 6: Effect of time to soursop fermentation**



After 60 hours of fermentation, we get the best soursop fermented juice.

**Effect of concentrated juice supplementation to soursop fermented juice quality**


**Fig. 7: Effect of concentrated juice supplementation to quality of the fermented juice**



We see that 30% supplementation of concentrated juice is appropriated to get the best fermented juice.

**Effect of syrup supplementation to soursop fermented juice quality**


**Fig. 8: Effect of syrup supplementation to soursop fermented juice quality**



With 8% syrup supplementation, we get the best soursop fermented juice.

**Effect of caramel supplementation to soursop fermented juice quality**


**Fig. 9: Effect of caramel supplementation to soursop fermented juice quality**




With 1.5% caramel addition, we get the best soursop fermented juice.

**Effect of the pasteurization temperature to the soursop fermented quality**

**Fig. 10: Effect of the pasteurization temperature to the soursop fermented quality**



At 80°C of pasteurization we get the best sourop fermented quality so we choose this value for application.

**Effect of the pasteurization time to the soursop fermented quality****Fig. 11: Effect of the pasteurization time to the soursop fermented juice quality**

Pasteurization which is conducted within 10 minutes will have the best soursop fermented juice quality.

**CONCLUSION**

Soursop also known as guanabana (*Annona muricata* L.) belonging to the family *Annonaceae* and indigenous one of the exotic fruits prized for its very pleasant, sub-acid, aromatic and juicy flesh which consists of edible white pulp and a core of indigestible black seeds. This study has shown that low alcoholic wine with appreciable qualities and immense acceptability can be produced from ripe soursop fruit.

**REFERENCES**

1. Abbo, E.S. Olurin, T. Odeyemi, G., Studies on the storage stability of soursop (*Annona muricata* L.) juice. *Afr J Biotechnol.* **5**: 108–112 (2006)
2. Asprey, G.F. Thornton, P., Medicinal plants of Jamaica. *West Indian Med.* **4**: 69–92 (1995)
3. Bates, R.P. Moris, J.R. Candall, P.G., Principles and practices of small and medium scale fruit juice processing. *FAO United Nations Newsletter.* Rome, Italy: United Nations; 2001.
4. Imade, E.E. Ikenebomeh, M.J. Obayagbona, O. N. and Igiehon, O.N., Evaluation of changes in the microbial profile, physico-chemical and nutritional attributes during the bioconversion of soursop (*Annona muricata*) must to wine. *Nig J. Biotech.* **25**: 1-11 (2013)
5. E. Mbaeyi-Nwaoha, C. N. Ajumobi, Production and microbial evaluation of table wine from tamarind (*Tamarindus indica*) and soursop (*Annona muricata*). *Journal of Food Science and Technology*, **52(1)**:105-116 (2015)
6. Nwachukwu, E. and Ezeigbo, C. G., Changes in the microbial population of pasteurized soursop juice treated with benzoate and lime during storage. *African Journal of Microbiology Research*, **7 (31)**: 3992-3995 (2013)
7. Ogbemor, C. Akpoveta, V.O. Osakwe, S.A. Medjor, W.O., Fermentation of soursop using *Saccharomyces Cerevisiae*: a kinetic evaluation. *Chemistry and Materials Research*, **6(9)**: 60-64 (2014)
8. Raphael N Okigbo and Omokaro Obire, Mycoflora and production of wine from fruits of soursop (*Annona Muricata* L.). *International Journal of Wine Research* **1**: 1–9 (2009)
9. Sunday, P. Ukwo, Chidi, F. Ezeama, Studies on proliferation of acetic acid bacteria during soursop juice fermentation. *Internet Journal of Food Safety*, **13**: 345-350 (2011)