

Oilseeds Potential of *Zanthoxylum zanthoxyloides* (Lam.) Zepernick and Timler

Lassina Ouattara^{1*}, Paulin Ouoba², Philippe-Pierre Robichaud³ and Marc E. Surette³

¹Université Polytechnique de Bobo-Dioulasso, Unité de Formation et de Recherche en Sciences et Techniques (UFR-ST), Département de Génie-Biologie. 01 BP 1091 Bobo-Dioulasso 01. Burkina Faso

²Université Polytechnique de Bobo-Dioulasso, Unité de Formation et de Recherche en Sciences et Techniques (UFR-ST), Département de Sciences Biologiques. 01 BP 1091 Bobo-Dioulasso 01. Burkina Faso

³Université de Moncton, Département de Chimie et Biochimie, Moncton, NB, E1A 3E9 Canada

*Corresponding Author E-mail: lasouatt@yahoo.fr

Received: 20.10.2016 | Revised: 28.10.2016 | Accepted: 29.10.2016

ABSTRACT

The aim of this prospective study was to assess firstly oil potential of *Zanthoxylum zanthoxyloides* fruit and then the fatty acid profile. After extracting the oil from the seeds of this plant by the soxhlet method using petroleum ether as solvent, we proceeded to analyze the profile of fatty acids by gas chromatography with derivatization. A very interesting oil yield of $41.8 \pm 4.0\%$ was measured and highlighted a total of thirteen fatty acids with six saturated fatty acids and seven unsaturated fatty acids at varying proportions. This oil contained a high proportion of unsaturated fatty acids (72.4%) and the major unsaturated fatty acids are oleic acid ($46.4 \pm 12.7\%$), linoleic acid ($12.8 \pm 3.5\%$) and alpha-linolenic acid ($11.7 \pm 3.2\%$). This oil has a particular fatty acid profile due to the presence of three essential fatty acids namely linoleic acid ($\omega 6$) and alpha-linolenic acid ($\omega 3$) in large quantities and very small quantities of gamma linolenic acid ($\omega 6$).

Key words: *Zanthoxylum zanthoxyloides*, oilseed, fatty acids.

INTRODUCTION

In sub-Saharan Africa many plant oils are nowadays used in industry^{6,9}. In Burkina Faso in 2014, production of groundnut, sesame, soy and cotton seed were respectively 335,223; 321,837; 15,055 and 894,982 metric tons⁷. These speculations are the main sources of edible extracting oil in the country. Despite this production, Burkina Faso remains an oil importing country¹², hence the need to diversify sources of production. Many species

found in African flora are still underutilized, even unrecognized by populations¹³. Yet these species are potential sources of new edible oils. This is the case of *Zanthoxylum zanthoxyloides* (Lam.) Zepernick and Timler. No study made on the seeds of this species has mentioned it as the oleaginous species. *Zanthoxylum zanthoxyloides* belongs to the family of Rutaceae. It is a shrub or small tree, prickly reaching 12 meters high.

Cite this article: Ouattara, L., Ouoba, P., Robichaud, P.P. and Surette, M.E., Oilseeds Potential of *Zanthoxylum zanthoxyloides* (Lam.) Zepernick and Timler, *Int. J. Pure App. Biosci.* 4(5): 23-28 (2016). doi: <http://dx.doi.org/10.18782/2320-7051.2394>

The bark is gray to beige, rough, with thin vertical cracks and often woody protrusions wearing spurs. The leaves are alternate, smooth, imparipinnately compound, 5 to 11 leaflets opposite or alternate and with many glandular points. The inflorescence is a terminal or axillary panicle lose 5-25 cm long, ovoid fruits are dehiscent follicles, 5-6 mm in diameter, red and shiny black seeds³. This plant is found in the tropical zone of West and Central Africa². In Burkina Faso, this species is very common in some Southwest flora¹⁴.

The goal of this work is to highlight the oleaginous potential of the seeds of this plant

and to determine the fatty acid profile of its oil. This work is in relationship with the characterization and the valuation of the Burkina Faso plant resources.

MATERIALS AND METHODS

Plant material

Zanthoxylum zanthoxyloides fruits were harvested in the natural forests of Dindérosso and Kua, two places located around the city of Bobo-Dioulasso. In these places, flowering of *Zanthoxylum zanthoxyloides* begins in July and then fruits (Figure 1) are available from October until March.



Fig.1: Ripe fruits of *Zanthoxylum zanthoxyloides* (Source: Ouoba, 2006)

Seed harvesting

The fruits were harvested during November and December. In the laboratory the fruits were dried and stored away from light. When the follicles are dry they open to release the seeds.

Seed oil extraction

Dried seeds of *Zanthoxylum zanthoxyloides* were milled with a crusher type Retsch GM-Copyright © October, 2016; IJPAB

200. Homogenates obtained were used for extraction of the fat matter. The extraction was done using a soxhlet extractor according to the NF EN ISO 659¹⁰. For this, the cartridges (HM1004-33x94mm) were filled with 200g grams of the homogenate and placed in the extractor column. As extracting solvent, petroleum ether was used (Carlo Erba SDS-REACTIVE), an apolar solvent. After 4 hours,

the extract (mixture of ether and fat) was evaporated (Rotavapor R-200 BÜCHI) to remove the solvent. The experiment was repeated three times. The fat, stripped of solvent, was recovered and the average extraction yield calculated using the following formula:

$(m / M) \times 100$, where **m** is the oil mass and **M** the mass of mash. The oil obtained was stored at 4°C before analysis of the profile of fatty acids by gas chromatography.

Fatty acids profile analysis

The crude oil obtained by soxhlet extraction was analyzed by gas chromatography derivatization¹⁵. The method can be briefly described as follows: We mixed in a glass tube 2 ml of methanol, 0.9 ml of chloroform, 20 microliters of acetic acid, 800 microlitres of MilliQ water, 20 microliters of the crude oil of *Zanthoxylum zanthoxyloides* and 100 microliters of methanol containing the internal standard (C17:0). After 15 min incubation at room temperature, 2 ml of chloroform and 1 ml of MilliQ water were added. The tubes were gently inverted twice and centrifuged 5 min at 1000 rpm to separate the two phases. The lower phase (chloroform) was transferred to another glass tube and a second extraction was done on the first tube with 2 ml of chloroform and the two lower phases were combined. The lipids were dried and hydrolyzed with 400 microliters of KOH (0.5 M) at 100° C for 15 min. Then the fatty acids were transmethylated with 1 ml of boron trifluoride (14% in methanol) at 100° C for 10 min. The FAMES were then extracted by adding 2 ml of hexane and 1.6 ml of HCl (0.125 M). After homogenization with a vortex, the upper phase was transferred to a GC vial and analyzed by gas chromatography using a Thermo Trace gas chromatograph (Thermo Electron Corporation, Mississauga, ON, Canada). The volume injected was 1

microliter. The injection was made in split mode 1/40 onto a Trace-FAME column (Thermo) with flame ionization type detector. The fatty acids were identified by comparing their retention times with those of authentic standards, and a standard curve was used for quantification¹⁵.

Data analysis

Microsoft Excel was used as data analysis software. The oil extraction yields and percentages of fatty acids are expressed as mean plus or minus standard deviation values.

RESULTS

Oil extracting yield

After performing the extraction three times a yield of $41.8 \pm 4.0\%$ was measured.

Fatty acid composition of the crude oil of the seeds of *Zanthoxylum zanthoxyloides*

We have noted in the crude oil, thirteen measurable fatty acids including six saturated fatty acids and seven unsaturated fatty acids (Figure 2 and Table 1). In terms of proportion, this oil is more concentrated in unsaturated fatty acid (72.35%) than saturated fatty acids (27.65%). The majority unsaturated fatty acids are: oleic acid ($46.42 \pm 12.65\%$), linolenic acid ($11.73 \pm 3.17\%$) and linoleic acid ($12.82 \pm 3.46\%$). The majority of saturated fatty acids are: palmitic acid ($23.86 \pm 6.40\%$) and stearic acid ($3.27 \pm 0.9\%$) also minority fatty acids in descending order are: vaccenic acid, palmitoleic acid, arachidic acid, lignoceric acid, behenic acid, myristic acid and finally gamma linolenic acid. This oil has a particular fatty acid profile, due to the presence of three essential fatty acids namely linoleic acid ($\omega 6$), linolenic acid ($\omega 3$) in large quantities and gamma linolenic acid ($\omega 6$) in small quantities. Monounsaturated fatty acids represent 47.73% of total fatty acids against 24.62% for polyunsaturated fatty acids.




Fig.2: Chromatogram of *Zanthoxylum zanthoxyloides* seeds oil giving retention time and relative abundance of fatty acids

Table 1: Fatty acids of *Zanthoxylum zanthoxyloides* seeds oil

Fatty acid		Percentage (%)	Retention time (min)
C14:0	Myristic acid	0.07 ± 0.02	3.51
C14:1	Myristoleic acid	0.00	-
C16:0	Palmitic acid	23.86 ± 6.4	4.63
C16:1	Palmitoleic acid	0.34 ± 0.09	4.97
C18:0	Stearic acid	3.27 ± 0.9	6.31
C18:1_n-9	Oleic acid	46.42 ± 12.65	6.74
C18:1_n-7	Vaccenic acid	0.53 ± 0.14	6.82
C18:2_n-6	Linoleic acid	12.82 ± 3.46	7.47
C18:3_n-6	Gamma linolenic acid	0.06 ± 0.02	8.10
C18:3_n-3	Alpha linolenic acid	11.73 ± 3.17	8.49
C20:0	Arachidic acid	0.28 ± 0.08	8.78
C18:4_n-3	Stearidonic acid	0.00	-
C20:1	11-eicosenoic acid	0.44 ± 0.10	9.34
C20:2_n-6	11,14-eicosadienoic acid	0.00	-
C20:3_n-6	Gamma-linolenic acid	0.00	-
C20:4_n-6	arachidonic acid	0.00	-
C20:3_n-3	11,14,17-eicosatrienoic acid	0.00	-
C22:0	Behenic acid	0.08 ± 0.02	12.17
C22:1_n-9	Erucic acid	0.00	-
C20:5_n-3	Eicosapentaenoic acid	0.00	-
C22:4_n-6	Docosatetraenoic acid	0.00	-
C24:0	Lignoceric acid	0.09 ± 0.03	16.46
C24:1_n-9	Nervonic acid	0.00	-
C22:5_n-3	Docosapentaenoic acid	0.00	-
C22:6_n-3	Docosahexaenoic acid	0.00	-

DISCUSSION

After extracting the oil in triplicate, we obtained an average oil yield of $41.77 \pm 3.98\%$. This shows that this plant has an interesting oilseed potential. This yield is quite high compared to several common oilseeds as cotton and soya beans contain, respectively, 15-25% and 15-22% oil⁹.

The analysis of the profile of fatty acids of the crude oil has allowed us to highlight the presence of six saturated fatty acids and seven unsaturated fatty acids at various concentrations. This oil has a particular fatty acid profile, due to the presence of three essential fatty acids namely linoleic acid ($\omega 6$), linolenic acid ($\omega 3$) in large quantity and gamma linolenic acid ($\omega 6$) in small quantities.

In terms of comparison, peanut oil which is among the most commonly used oils in the region of western Burkina Faso, contains omega-9 and omega-6 fatty acids, which were also measured essential fatty acids in oil of *Zanthoxylum zanthoxyloides*. But peanut oil only contains traces of omega-3 fatty acids^{5,4}. Cotton oil is also poor in omega-3 fatty acids, while sesame oils contains both omega-6 and omega-3 fatty acids⁵.

In addition the ratio of the proportions of polyunsaturated fatty acids to saturated fatty acids was 0.89, which is slightly above 0.7 considered by some authors as ideal for cooking oil⁸. The ratio of omega-6/omega-3 fatty acids is 1.1 and according to Lambert⁵, the maximum ratio of omega-6/omega-3 for edible oil should be 5 thus this ratio for *Zanthoxylum zanthoxyloides* oil is well below this maximum value. This ratio is also lower compared to thirteen other edible oils such as grapeseed oil, natural sunflower, cotton, peanut, oleic sunflower, corn germ, safflower, soybean, margarine, primrose, olive, walnut, hemp and rapeseed⁵.

Oleic acid (monounsaturated fatty acid) is the major fatty acid measured in *Zanthoxylum zanthoxyloides* oil with a proportion of $46.42 \pm 12.65\%$ of the total fatty acids, that approximates that of peanut oil⁵. Oleic acid is cited by some authors to have beneficial

properties in relation to an extended shelf life and high thermal stability⁶ and its elevated concentration in *Zanthoxylum zanthoxyloides* oil is a desirable trait.

The proportion of the unsaturated acid is 72.35% of the total fatty acids; which is close to the proportion of unsaturated fatty acids in cotton seed oil⁵, with the saturated palmitic acid making up most of the rest of the fatty acid profile as it is the second most important fatty acid in terms of proportion, $23.86 \pm 6.40\%$ of total fatty acids. All these results support the use of this oil as food. In addition, the fruits of this species are used as spices in the sauces in some African countries such as Cameroon¹, therefore the seed oil is not the only edible portion of the fruit.

Conclusion and perspectives

This study allowed us to identify the oilseed potential of the seeds of *Zanthoxylum zanthoxyloides* and to specify the fatty acid composition of its oil. These seeds are a rich source of oil which makes up nearly 42% of the dry weight of the seeds. The oil has a particular fatty acid profile because of its high proportion of unsaturated fatty acids (72.35%) with the remarkable presence of oleic acid with a proportion of $46.42 \pm 12.65\%$ of the total of fatty acids, and a very desirable omega-6/omega-3 fatty acid ratio of 1.1. Having regard to the profile of these fatty acids, the oil could be used as an edible oil after further studies such as analysis of glycerides, phospholipids and dyes. A cytotoxicity study would also be warranted. Once these studies are successful, we will consider an extension of the production of this plant as an oilseed plant.

Acknowledgments

Department of Chemistry and Biochemistry of the University of Moncton (CANADA). Especially we want to thank the technical team from Marc Surette's laboratory for their contribution during the analysis of this oil. Marc Surette is the recipient of an Innovation Research Chair in Biosciences from the New Brunswick Innovation Foundation.

REFERENCES

1. Abdou Bouba, A., *Contribution à l'étude du développement d'un aliment fonctionnel à base d'épices du Cameroun : Caractérisation physico-chimique et fonctionnelle*. Thèse de Doctorat, de l'Université de N'Gaoundéré, Cameroun (2009).
2. Arbonnier, M., *Arbres, arbustes et lianes des zones sèches d'Afrique de l'Ouest*. CIRAD-MNHN-UICN, (2002).
3. Berhaut, J., *Flore illustrée du Sénégal*. Ministère du Développement Rural et de l'Hydraulique, Direction des Eaux et Forêts, Dakar. Tome 1 (1971).
4. Carrin, M.E. and Carelli, A.A., Peanut oil: Compositional data. *European Journal of Lipid Science and Technology*. **112**: 697–707 (2010).
5. Lambert, J., *Les huiles végétales : 2000 plantes oléagineuses*. Institut Français des huiles végétales pures (2005).
6. Leray, C., *Les lipides dans le monde vivant : introduction à la lipidologie*. Edition Lavoisier (2010).
7. FAOSTAT., 2014: <http://faostat3.fao.org/download/Q/QC/> (consulted on april 2, 2016)
8. Fossati, P., Bases nutritionnelles de l'alimentation lipidique normale. *Thérapie.*, **51**: 615- 621 (2000).
9. Ribier, D. and Rouzière, A., *La transformation artisanale des plantes à huile : expériences et procédés*. Paris: GRET (Guide pratique: GRET) (1995).
10. Norme NF EN ISO 659 (2009).
11. Nzikou, J.M., Matos, L., Bouanga-Kalou, G., Ndangui, C.B., Pambou-Tobi N.P.G., Kimbonguila, A., Silou Th, Linder, M. and Desobry, S., Chemical Composition on the Seeds and Oil of Sesame (*Sesamum indicum* L.) Grown in Congo-Brazzaville. *Advance Journal of Food Science and Technology*, **1(1)**: 6-11(2009).
12. OCL-Oilseeds&fats and Crop and Lipids., Burkina Faso: Une approche innovante partagée par Agropol et la FAO. *EDP Sciences*, **21(1)**: N101:5 (2014).
13. Ouoba, P., Boussim, I. J., Lykke, A.M. et Guinko S., Le potentiel fruitier de la Forêt Classée de Niangoloko. *Fruits.*, **61(1)**: 71-81(2006).
14. Ouoba, P., *Flore et végétation de la Forêt Classée de Niangoloko*. Thèse de Doctorat, de l'Université de Ouagadougou, Burkina Faso (2006).
15. Surette, M.E., Koumenis, I.L., Edens, M.B., Tramposch, K.M. and Chilton, F.H., Inhibition of leukotriene synthesis, pharmacokinetics, and tolerability of a novel dietary fatty acid formulation in healthy adult subjects. *Clin Ther.*, **25(3)**: 948-971(2003).