

Carbon Sequestration under Different Cropping Systems with Different Depth and Its Impact on Climate Change

Bhavya V. P. *, Anil Kumar S., Alur, A., Shivakumar, K. M. and Shivanna, M.

Department of Soil Science and Agricultural Chemistry, College of Horticulture, Bengaluru-560 065, Karnataka

*Corresponding Author E-mail: ammushetty1993@gmail.com

Received: 21.12.2017 | Revised: 26.01.2018 | Accepted: 3.02.2018

ABSTRACT

The main objective of this study was to assess the carbon stocks and carbon dioxide sequestration under different land use systems within the same locality with twenty year old cultivation. The soil samples were collected at a depth of 0-15cm, 15-30cm, 30-50 cm and 50-100 cm from a mango, cashew, rose, vegetable and medicinal and aromatic cropping systems. The study showed that, the magnitude of carbon sequestration is more under mango orchard followed by cashew orchard than annuals crops like rose, medicinal and aromatic and vegetable block. The carbon dioxide sequestration was significantly greater under the perennial crops as compared to annual crops. It was observed that perennial horticulture crops increases the soil organic carbon (SOC) and carbon dioxide storage than annual crops and reduce the carbon emissions to the atmosphere which helps to mitigate the global warming.

Key words: Carbon storage, Carbon dioxide sequestration, Climate change, Land use systems

INTRODUCTION

Carbon sequestration by terrestrial ecosystem is the net removal of carbon dioxide (CO₂) gas from the atmosphere or the avoidance of its emissions from terrestrial ecosystems into the atmosphere. The removal process include: CO₂ uptake from the atmosphere by green plants through photosynthesis and the carbon stored as plant biomass (in the trunks, branches, leaves and roots of the plants) and organic matter in the soil. Soil carbon sequestration involves adding the maximum possible organic carbon to the soil. The carbon dioxide gas is a major cause of the atmospheric greenhouse effect and hence can influence the global climate. However, many other gases such as methane (CH₄), ozone (O₃) and nitrous

oxide (N₂O) influence the global climate. Carbon sequestration studies have gained momentum in the recent decade and the amount of carbon stored in a system is a good measure of its sustainability. Most of the factors affecting carbon sequestration are affected by land management practices. The distortion in the global carbon balance through human activities is due to the burning of fossil fuel and agriculture and land use change (33%). Actions to sequester carbon in the soils will generally increase the organic matter content of soils. Soil organic carbon, as reported by Bationo is simultaneously a source and sinks for nutrients and plays a vital role in soil fertility maintenance.

Cite this article: Bhavya, V.P., Kumar, S.A., Alur, A., Shivakumar, KM. and Shivanna, M., Carbon Sequestration under Different Cropping Systems with Different Depth and Its Impact on Climate Change, *Int. J. Pure App. Biosci.* 6(1): 1612-1616 (2018). doi: <http://dx.doi.org/10.18782/2320-7051.6114>

Growing of perennial horticulture crops is one of the strategies to improve soil conditions which would result in enhancing soil attributes and contributing to the good soil health and helps to sequester more organic carbon and carbon dioxide in soil as compared to annual crops. Although the benefits of perennial horticulture crops on soil in improving its chemical properties have been well known, but the present study is a modest attempt to assess the carbon storage and carbon dioxide sequestration under diverse horticultural cropping systems within the same locality of four year old cultivation *i.e.*, annuals *vis-a-vis* perennial crops. The hypothesis of the study was that, land use systems sequester SOC and CO₂ differently.

MATERIAL AND METHODS

The present investigation was carried out at division of horticulture, UAS Campus, GKVK, Bengaluru, Karnataka during 2015. The soil of the experimental site was sandy loam and classified as fine, mixed isothermic Kandic Paleustalf of Vijayapura soil series. Cultivation of horticultural crops *viz.*, mango, cashew, rose, vegetables and medicinal and aromatic crops were selected for the experiment. The experimental plots were permanently laid out for specified crops. In each sampling site, soil samples were collected from four different depths *i.e.*, 0-15, 15-30, 30-50, and 50-100 cm. The study was carried out in a Randomized Complete Block Design (RCBD). Soil samples were processed by

drying under the shade, powdering with a clean wooden mallet and passing through a 2 mm sieve. The processed samples were stored in polyethylene bags for analysis in the laboratory. For determination of organic carbon, a small volume of stored sample was further ground and passed through a 0.2 mm sieve before analysis and soil organic carbon (SOC) was determined by the modified Walkley and Black dichromate digestion method.

Calculation of the organic carbon sequestered (stored)

The SOC stored was calculated by using the Donovan formula.

$$CT = CF \times D \times V$$

Where:

CT is the total carbon stored per area in Mg ha⁻¹

CF is the fraction SOC (percentage carbon divided by hundred), D is the soil bulk density and V is the volume of the soil in cubic meters (which is = Depth of soil x area of soil)

Conversion of soil organic carbon to CO₂

The final results were multiplied by a factor of 3.67 (*i.e.* the molecular mass of CO₂/ atomic mass of C) to convert the total carbon stored to carbon dioxide.

RESULTS AND DISCUSSIONS

Soil organic carbon under different land use systems

Organic carbon content at different depths as influenced by different horticulture crops is presented in figure 1.




Fig. 1: MO- Mango Orchard, VB- Vegetable Block, CO- Cashew Orchard, MAP's- Medicinal and Aromatic Plants, RB- Rose Block

The organic carbon content of soil was higher in the surface layer of the soil *i.e.*, 0-15 cm as compared to other subsurface soil depth in all the cropping systems and the organic carbon content decreased with increase in soil depth. The carbon content differed significantly with different cropping system, the mango orchard had higher organic carbon content *i.e.*, 7500.00, 7334.00, 7016.00, and 6743.00 mg kg⁻¹ at 0-15 cm, 15-30 cm, 30-50 cm and 50-100 cm depths respectively which was followed by cashew orchard whereas, the medicinal and aromatic block had lowest organic carbon 4521.00, 4311.00, 4008.00 and 3786.00 mg kg⁻¹ at 0-15 cm, 15-30 cm, 30-50 cm and 50-100 cm depths respectively. This is due to the continuous addition of organic matter by continuous falling of leaves in perennials crops like mango and cashew orchard leads to accumulate more organic matter to the soil. Since organic manures are incorporated in to the surface and a major

portion of the left over residues of shallow rooted crops usually accumulate in the top few centimetre of the soil layers, there was possibility for a relatively greater accumulation of organic carbon in 0-15 cm soil as compared to the soils of lower layer. Similar results were obtained by Manjaiah *et al*⁴. Increase in soil carbon due to continuous addition of bio mass through leaf and roots have been reported by Geo Jose² in case of field crops and by Krishnappa Naik *et al*³.

Bulk density under different land use systems Bulk density at different depths as influenced by different horticulture crops is presented in figure 2. Bulk density of the soil changed significantly over years of cultivation of perennial and annual crops. It was found that the higher bulk density values were recorded in the subsurface layers of the soil as compared to surface layers and it was observed that bulk density increased with increase in soil depth.




Fig. 2: MO- Mango Orchard, VB- Vegetable Block, CO- Cashew Orchard, MAP's- Medicinal and Aromatic Plant, RB- Rose Block

The annual crops like, medicinal and aromatic block (1.56 Mg cm⁻³) had higher bulk density followed by vegetable block (1.55 Mg cm⁻³) and significantly lowest values were recorded under the mango orchard (1.50 Mg cm⁻³) at 50-100cm soil depth. This is due to the continuous addition of organic matter with minimum tillage treatments under perennial crops reduces the breakdown of soil aggregates which helps to reduce the bulk

density. However, in case of annual crops due to continuous cultivation, the soil aggregates are broken down into small soil particles, as the size of the soil particles decreases the bulk density increases.

Soil carbon storage/stocks under different land use systems

Soil carbon storage at different depths as influenced by different horticulture crops is presented in Table 1. The carbon stocks

differed at different depth. The higher carbon stocks was observed in the subsurface layer *i.e.*, 50-100 cm and lowest was observed in 0-15 cm, 15-30 cm and 30-50 cm. The carbon stocks were increased with increase soil depth. Among different cropping systems, the mango orchard had maximum carbon stocks (5057.25 Mg ha⁻¹) which is on par with the cashew orchard (4119.28 Mg ha⁻¹) and significantly

lowest values was observed under the medicinal and aromatic block (3067.74 Mg ha⁻¹). In one meter soil depth, the mango orchard sequestered maximum carbon content (10342.75Mg ha⁻¹) which was followed by cashew orchard (8783.35Mg ha⁻¹) and lowest organic carbon was sequestered under the medicinal and aromatic block (6364.03Mg ha⁻¹).

Table 1: Soil carbon storage at different depths as influenced by different horticulture crops

Horticulture land use system	> 20 year old cultivation				
	Carbon stocks (Mg ha ⁻¹)				
	0-15 cm	15-30 cm	30-50 cm	50-100 cm	Total (1 m depth)
Mango orchard	1597.50	1584.14	2103.86	5057.25	10342.75
Cashew orchard	1428.09	1412.58	1823.4	4119.28	8783.35
Rose block	1134.30	1097.55	1454.94	3470.04	7156.83
Vegetable block	1102.60	1043.70	1366.90	3205.40	6718.60
Medicinal and aromatic block	1037.56	1016.25	1242.48	3067.74	6364.03
SEm ±	70.56	68.45	88.45	210.45	431.45
CD at 5%	210.11	205.45	266.47	630.15	1315.56

The carbon stocks were higher under the perennial horticulture crops as compared to annual crops. This is due to the continuous addition of organic matter under perennial crops which leads to continuous decomposition of the accumulated organic matter under the system. The thick litter also reduced the amount of CO₂ gas emitted into the atmosphere. The implication is that, crop residue application as surface mulch could play an important role in the maintenance of soil organic carbon levels and productivity. Continuous cultivation of the annual crops led to a decrease in the amount of organic carbon

stored in the soil as compared to the perennial crops. In horticultural systems, because the economic parts of the plants are harvested, only a small percentage of the production becomes available for incorporation into soil to enhance the organic carbon pool. All the aboveground production may be harvested, leaving only the root biomass. The low SOC recorded under the annual crops could be attributed to the little organic inputs from its component crops.

Carbon dioxide sequestration under different land use systems

Table 2: Carbon dioxide sequestration at different depths as influenced by different horticulture crops

Horticulture land use system	4 year old cultivation			
	CO ₂ sequestration (Mg ha ⁻¹)			
	0-15 cm	15-30 cm	30-50 cm	50-100 cm
Mango orchard	5862.82	5813.79	7721.16	18560.10
Cashew orchard	5241.09	5184.16	6691.87	15117.75
Rose block	5241.09	4028.00	5339.62	12735.04
Vegetable block	4046.54	3830.37	5016.52	11763.81
Medicinal and aromatic block	3807.84	3729.63	4559.90	11258.60
SEm ±	268.55	250.45	325.45	731.45
CD at 5%	806.45	752.25	977.45	2314.45

The carbon dioxide sequestration was differed at different depth. The highest sequestered carbon dioxide was observed in the subsurface layer *i.e.*, 50-100 cm and lowest was observed in 0-15 cm, 15-30 cm and 30-50 cm soil depths. The carbon dioxide sequestration was increased with increase in soil depths. Among different cropping systems, the mango orchard had maximum carbon dioxide sequestration (18560.10Mg ha⁻¹) which is on par with the cashew orchard (15117.75Mg ha⁻¹) and lowest was observed under the medicinal and aromatic block (11258.60Mg ha⁻¹) (Table 2). The CO₂ sequestration is less under the annual cropping systems as compared to the perennial crops. This is due to the land use changes and soil degradation processes as well as the rapid decomposition of organic matter in cultivated soils were the major cause for the release of CO₂ from the systems, as the land use systems that added more residues recorded the less emissions of CO₂. Cultivation of annual crops reduces the carbon pools and increase CO₂ emissions.

CONCLUSIONS

The study showed that the land use systems that sequestered more carbon and less CO₂ emission were ranked as: mango > cashew > rose > vegetable > medicinal and aromatic plants. The land use systems *i.e.*, perennial horticulture crops like mango and cashew orchard that added more residues recorded less emission of CO₂ and finally, it was observed

that perennial crops increase the SOC content than annual crops. Soil C sequestration is a multiple purpose strategy, it restores degraded soils, enhances the land productivity, improves the diversity, protects the environment and reduces the enrichment of atmospheric CO₂ and mitigates climate change.

REFERENCES

1. Donovan, P., Measuring soil carbon change: A flexible, practical, local method. pp 1-56 (2013).
2. Geo Jose, Organic carbon distribution in soils under long- term rice cropping systems in coastal and southern dry zone of Karnataka, *M.Sc (Argi.) Thesis*, Uni. Agric. Sci., Bengaluru (2006).
3. Krishnappa Naik, C. S., Haricharan, B. R., Jayarama. And Surendra Swamy, H. B., Coffee soils of Chikmagalur district. *Indian Coffee.*, **52(10)**: 3-9 (1988).
4. Manjaiah, K. M., Voroney, R.P., And Sen, U., Soil organic carbon stocks, storage profile and microbial biomass under different crop management systems in a tropical agricultural ecosystem. *Biol. Fertil. Soils.*, **31**: 273-278 (2000).
5. Walkey, A. J. And Black, C. A., An extraction method of degtjar method for determining soil organic matter and proposed modification chromic acid titration method. *J Soil Sci.*, **37**: 29-38 (1934).