

## Molecular Detection of *Klebsiella* spp. from Poultry Meat

Kiran S. Modh<sup>1\*</sup>, B. C. Parmar<sup>2</sup>, J. H. Chaudhary<sup>3</sup>, B. B. Bhanderi<sup>4</sup>, J. B. Nayak<sup>5</sup>,  
Sonali Thakur<sup>6</sup>, Z. B. Pargi<sup>7</sup> and N. M. Patel<sup>8</sup>

<sup>1,2,3,5,6,7,8</sup>Department of Veterinary Public Health and Epidemiology,

<sup>4</sup>Department of Veterinary Microbiology,

College of Veterinary Science and Animal Husbandry, AAU, Anand, Gujarat, India

\*Corresponding Author E-mail: [modikiran86@gmail.com](mailto:modikiran86@gmail.com)

Received: 22.12.2020 | Revised: 28.01.2021 | Accepted: 6.02.2021

### ABSTRACT

The Poultry are quite often implicated as a major reservoir of human enteric pathogens, and several human infections were traced after consumption of food products of animal origin. Commensal and opportunistic Enterobacteriaceae present in the poultry gut can be indirectly transmitted to humans through the food chain. In this study a total 150 raw Poultry meat samples (poultry muscle) was collected in sterile test screw capped vials and immediately transferred to the laboratory at 4° C for processing and bacteriological investigation which include the morphological characterisation, culture characterisation, Biochemical test and genotypic confirmation. 17 (11.33%) samples out of 150 raw chicken samples were positive for *Klebsiella* species.

**Keywords:** *Klebsiella*, Food borne disease, Raw Poultry Meat.

### INTRODUCTION

The Poultry sector in India is valued at about Rs. 80,000 crores (2015-16) broadly divided into two sub-sectors-one with a highly organized commercial sector with about 80% (Rs. 64,000 crore) of the total market share and the other being unorganized with about 20% of the total market share of Rs. 16,000 Crore (National Action Plan for Egg & Poultry, 2019). Poultry are quite often implicated as a major reservoir of human enteric pathogens, and several human infections were traced after consumption of food products of animal origin. Most

commonly Enterobacteriaceae found in the poultry gut can be indirectly transmitted to humans through the food chain and *Klebsiella* spp. is one of the most common infectious disease affecting chicks causing great economic losses. (Paterson & Bonomo, 2005).

The isolation, identification and characterization of *Klebsiella* spp. in chicken have been accomplished from the retail Poultry meat market in Anand district. The genus *Klebsiella* was named by Trevisan (1885) to honour the clinical microbiologist Edwin Klebs.

**Cite this article:** Modh, K. S., Parmar, B. C., Chaudhary, J. H., Bhanderi, B. B., Nayak, J. B., Thakur, S., Pargi, Z. B., & Patel, N. M. (2021). Molecular Detection of *Klebsiella* spp. from Poultry Meat, *Ind. J. Pure App. Biosci.* 9(1), 511-518. doi: <http://dx.doi.org/10.18782/2582-2845.8566>

The first *Klebsiella* species ever described was a capsulated bacillus from patients with Rhinoschleoma (Brisse et al., 2009). *Klebsiella* species are Gram-negative, non-motile, usually encapsulated rod-shaped bacteria belonging to the family Enterobacteriaceae. *Klebsiella* species often occur in mucoid colonies. The genus consists of 77 capsular antigens (K antigens), leading to different serogroups (Janda et al., 2007).

Biochemically *Klebsiella* positive reaction for catalase test, Voges Proskauer, Urease and Citrate test, meanwhile were negative for Oxidase, Indole and Methyl red tests. Cultivation of isolate on Mac-Conkey agar gives lactose fermenting colonies, more or less dome shaped. Three major species of *Klebsiella* viz. *Klebsiella pneumoniae*, *Klebsiella oxytoca* and *R. ornithinolytica* are ubiquitous in nature and are often detected in foods, sewage, soil, plants and the gastrointestinal tracts of animals (Sękowska, 2017). *K. pneumoniae* has been isolated from meats, vegetables and pastries (Babu et al., 2013).

*Klebsiella pneumoniae* is widely distributed in the gastrointestinal, urinary, and respiratory tracts of healthy people. It causes opportunistic infections mainly nosocomial infections; it is a common hospital acquired pathogen causing severe respiratory infections such as pneumonia. (Sękowska, 2017). Phenotypic-based methods for identification of *Klebsiella pneumoniae* are time consuming with low sensitivity hence; the various molecular-based approaches were used to identify *Klebsiella pneumoniae*. PCR has been successfully identified among these methods as a valuable method that provides rapid, sensitive and accurate.

## MATERIALS AND METHODS

This research was conducted at the Department of Veterinary Public Health and Epidemiology, College of Veterinary Science and Animal Husbandry, Anand Agricultural University, Anand. A total 150 raw Poultry meat samples (poultry muscle) were collected from retail meat market in and around Anand

under aseptic precautions in sterile screw lid sample vials and immediately transferred to the laboratory at 4°C for further processing and microbiological analysis.

### Isolation & identification

Sample was inoculated into 5 ml of nutrient broth for bacterial growth at 24 hours at 37°C. The growth in Nutrient broth was transferred to Mac-Conkey agar and again incubated at 37°C for 24 hours for selective growth of Gram negative bacteria, for specific isolation of *Klebsiella* spp., the pink, mucoid coloured colonies was picked up for further overnight incubation in HiCrome *Klebsiella* Selective Agar base supplemented with *Klebsiella* Selective Agar supplement (Carbenicillin 25mg/500 ml) at 37°C. Magenta coloured individual colonies was picked up as instructed by manufacturer and pure cultured in Nutrient broth and simultaneously streaked in MHA (Mueller-Hintonagar) plates for further study. All the pure cultures were kept at 4°C for further use.

**Morphological characterization:** Glass slides were stained with Gram's stain and examined microscopically for morphological characteristics of the isolates.

**Cultural characteristics:** The colonial morphology onto Mac-Conkey's agar was studied.

**Biochemical characterization:** Catalase test, Oxidase test, Indol test, Methyl red test, Vogasproskauer test, Citrate utilization test and Urease test, Triple sugar iron test were performed.

**Genotypic identification:** Identification of *Klebsiella* genus by *Klebsiella* specific gene (GyrA gene) and *Klebsiella pneumoniae* species identified by KP-16S (NM3) gene using polymerase chain reaction (PCR).

**DNA extraction for *Klebsiella* Genus and *Klebsiella pneumoniae* isolates:** The DNA from *Klebsiella* Genus was extracted by boiling method. A loopful of pure culture was suspended in 100 µl of DNase and RNase free milliQ water in a sterilized micro centrifuge tube. The suspension was vortexed and then heated at 95°C for 10 mins in thermal cyclor. This was then centrifuged at 10000 rpm

for 6 mins so that the cell debris settle down. The upper aqueous phase was transferred to another PCR tube and this was used as a DNA template for PCR.

**Oligonucleotide primers set GyrA:** Template DNA (5µl), forward and reverse primers (1µl), 12.5µl of master mix (2 times) (Fermentas, India) and 5.5µl of DNase free water (Fermentas, India) in a total volume of 25 µl. The melting temperature  $T_m$  of each oligonucleotide using the formula  $T_m = 4(G+C) + 2(A+T)$ , where G, C, A and T indicate the number of corresponding nucleotides in the oligonucleotide. Razmyar et al. (2015).

**Oligonucleotide primers set KP-16S (NM3) for *Klebsiella pneumoniae*:** Template DNA (5µl), forward and reverse primers (1µl), 12.5µl of master mix (2 times) (Fermentas, India) and 5.5µl of DNase free water (Fermentas, India) in a total volume of 25 µl. The melting temperature  $T_m$  of each oligonucleotide using the formula  $T_m = 4(G+C) + 2(A+T)$ , where G, C, A and T indicate the number of corresponding nucleotides in the oligonucleotide. Aurna et al. (2017).

#### PCR protocol:

- 1) The reaction was included in a total volume of 25µL in 0.5 ml Eppendorf tube containing template DNA (5µl), forward and reverse primers (1µl), 12.5µl of master mix (2 times) and 5.5µl of DNase free water.
- 2) The tubes were placed in the thermal cycler previously programmed.
- 3) At the end of cycling the tubes were stored at - 20°C until needed for electrophoresis.

#### Cyclic condition for detection of GyrA gene:

The thermal cycler was programmed as follows: (i) one cycle for 5 minutes at 94°C for initial denature the template DNA followed by (ii) 30 cycles of denaturation, annealing and extension at 94°C for 45 sec. and, 55° for 30 sec and 72° C for 45 sec. The 30 cycles were followed by a final cycle of extension at 72°C for 4 minutes to ensure that the entire PCR product was double strand DNA.

#### Cyclic condition for detection of gene KP-16S (NM3):

The thermal cycler was programmed as follows:(i) one cycle for 3 minutes at 94°C for initial denature the template DNA followed by (ii) 35 cycles of denaturation, annealing and extension at 94°C for 30 sec. and, 58° for 30 sec and 72° C for 30 sec. The 35 cycles were followed by a final cycle of extension at 72°C for 10 minutes to ensure that the entire PCR product was double strand DNA.

## RESULTS

#### Results of poultry chicken:

A total 150 raw Poultry meat samples (poultry muscle) revealed 17 isolates of *Klebsiella* with percentage of 11.33%. (Table 1).

#### Identification of suspected isolates:

**Morphological identification of *Klebsiella*:** Gram negative bacilli, non-spore forming, capsulated, non-motile and arranged singly. (Fig 1).

**Culture character:** on mac-Conkey Rose Pink Mucoid coloured colonies. (Fig 2) On selective media of *Klebsiella* selective agar base supplemented with Carbenicillin (25 mg/500 ml) produce magenta coloured colonies. (Fig3).

#### Biochemical identification:

Gave positive reaction for catalase test, Vogasproskauer test, citrate test and urease test. Meanwhile the isolates were negative for indole, oxidase and methyl red tests, and Triple sugar iron test produce acid slant and acid butt. (Table 1) (Fig 4) (Fig 5) (Fig 6).

#### Results of PCR:

Result of polymerase chain reaction for detection of *Klebsiella* genus specific GyrA gene were 12 isolates give positive results. (Table 2) (FIG 7).

Result of polymerase chain reaction for detection of *Klebsiella pneumoniae* by using KP-16S (NM3) gene were 9 isolates give positive results. (Table 3) (FIG 8).

## DISCUSSION

Results demonstrated by examination of 150 raw chicken samples revealed the percentage of *Klebsiella* infection among the examined

chicks was found to be 11.33%. The obtained results agree with that of Guo et al. (2016) 11.4%, Veeraselvam et al. (2019) 15.13%, Wu et al. (2016) 17.0% and are differed with the result received by Gundogan et al. (2011) 80%, Fielding et al. (2012) 40%, Mahanti et al. (2017) 65%, Eibach et al. (2018) 29%, Von Tippelkirch et al. (2018) 31.8% who recorded higher isolation rate of *Klebsiella* associated with heavy contamination, unhygienic processing of poultry meat. Also lower prevalence rate was recorded by Aly et al. (2014) 10.66%, Hayati et al. (2019) 9.21% and Hermana et al. (2020) 2.5%. Among the positive *Klebsiella* spp. 52.94% strains were confirmed of *Klebsiella pneumoniae* to be positive by PCR. The obtained results agree with that of Rajaa, et al. (2011) who recorded higher isolation rate of *Klebsiella pneumoniae* (40.4), Kurupati et al. (2004) 100%.

The selected isolation media used in this test was Mac-Conkeys agar media after incubation 24 hrs colonies appear rose pink mucoid

coloured colonies after overnight incubation at 37°C. This result agree with that of Aher et al. (2012). Total of 17 positive isolates were confirmed as *Klebsiella* spp. on the basis of standard biochemical tests. All the purified *Klebsiella* spp. isolates were exhibited standard biochemical results like catalase (+ve), oxidase (-ve), Indole - Methyl red - Voges Proskauer – (Citrate utilization on Simmon's citrate medium), (IMViC) (- or +, -, +, +) and H<sub>2</sub>S production on TSI agar and this result agree with Alves et al. (2006). In the present study, GyrA (*Klebsiella* genus) and KP-16S (NM3) (*Klebsiella pneumoniae*) genes were detected by PCR using specific primer sequences which yielded product sizes of 441 bp and 657 bp respectively. Out of total 17 isolates, 12 isolates (70.58%) was positive for GyrA gene this result concised with result of Aly et al. (2014) and (Younis et al., 2016) and 9 isolates (52.94%) was positive for KP- 16S (NM3) gene this result concised with result of (Kurupati et al., 2004).

**Table 1: Prevalence of *Klebsiella* spp.**

Types of samples	Total No. of samples collected	No. of samples positive	Prevalence percentage
Poultry Muscle	150	17	11.33

**Table 2: Results of the biochemical tests performed for all *Klebsiella* spp. Isolates**


Sr.no	Biochemical tests performed for the confirmation of culturally isolated <i>Klebsiella</i> spp.	Results for all <i>Klebsiella</i> spp. isolates
1	Indol Production	-ve
2	Methyl-red test	-ve
3	Voges-Proskauer test	+ve
4	Citrate utilization on Simmon's citrate medium	+ve
5	Catalase test	+ve
6	Oxidase test	-ve
7	Urease test	+ve
8	Triple sugar ion test H <sub>2</sub> S production on TSI agar	Yellow slant, Red butt, H <sub>2</sub> S production

**Table 3: Detection of *Klebsiella* spp. by PCR**

Sr. No.	Name of Genes	No. of isolates found positive	Prevalence Percentage
1	<i>gyrA</i> (441 bp)	1 2	70.58 %

**Table 4: Molecular Detection of *Klebsiella pneumoniae* gene by PCR**

Sr. No.	Name of Genes	No. of isolates found positive	Prevalence Percentage
1	KP-16S (NM3) 657bp	9	52.94 %




**Figure 1: Gram negative, Non- motile, Encapsulated and Rod shaped bacterium**



**Figure 2: Rose pink Mucoïd coloured colonies on MacConkey agar**



**Figure 3: Magenta coloured colonies on *Klebsiella* selective agar**



**Figure 4:**

- 1) Indole negative
- 2) Methyl red negative
- 3) Voges-Proskauer positive
- 4) Citrate positive




Figure 5: Urease test



Figure 6: TSI test




Figure 7: PCR amplification of *gyrA*(441 bp) genes  
 Lane M: 100 – 600 bp plus DNA ladder (Qiajen, Germany)  
 Lane 1: Positive control (MTCC 7407 *Klebsiella Pneumoniae*)  
 Lane 2: 2-6: Samples positive for *gyrA* gene of *Klebsiella spp.*  
 Lane 3: 7: Samples negative for *gyrA* gene of *Klebsiella spp.*(Negative control)




Figure 8: PCR amplification of *KP-16S* (NM3) (657bp) genes Lane M: 100 bp plus DNA ladder (Genei, Bangalore)  
 Lane N: Negative control  
 Lane 1, 2, 5-10: Samples positive for *KP-16S* gene of *Klebsiella spp.*  
 Lane 3, 4: Samples negative for *KP-16S* gene of *Klebsiella spp.*  
 Lane 11: Positive control (MTCC 7407 *Klebsiella Pneumoniae*)

### CONCLUSIONS

Out of total 150 chicken samples (Raw chicken), 17 (11.33%) samples were found positive for *Klebsiella species* by cultural isolation. All the *Klebsiella* spp. isolates showed mucoid pink colonies on Mac-Conkey's Agar medium and were further processed in *Klebsiella* Selective Agar which showed colonies with Magenta colour and all the 17 *Klebsiella* spp. isolates exhibited similar IMViC pattern of - - + +, produced negative result for Indole and Methyl Red (MR) and positive for Vogasproskauer (VP), citrate utilization test. Out of 17 isolates investigated, *Klebsiella* spp. isolates were found a prevalence rate of 12(70.58%) gyrA (441bp) gene and 9(52.94%) KP-16S (NM3) (657bp) by PCR.

### REFERENCES

- Aher, K. R. (2012). Geochemistry and assessment of ground water quality for drinking and irrigation purposes: a case study of Sukhana river subbasin, district Aurangabad, Maharashtra, India. *International Journal of Recent Trends in Science and Technology*, 4(1), 45-49.
- Alves, M. S., da Silva Dias, R. C., de Castro, A. C. D., Riley, L. W., & Moreira, B. M. (2006). Identification of clinical isolates of indole-positive and indole-negative *Klebsiella* spp. *Journal of clinical microbiology*, 44(10), 3640-3646.
- Aly, M. M., Khalil, S., & Metwaly, A. (2014). Isolation and Molecular Identification of *Klebsiella* Microbe Isolated from Chicks. *Alexandria Journal for Veterinary Sciences*, 43(1), 97-103.
- Aurna, S. T. (2017). *Rapid identification of Klebsiella pneumoniae* using PCR based method targeting 16S *rRNA* gene (Doctoral dissertation, BRAC University).
- Babu, L., Reddy, P., Murali, H. S., & Batra, H. V. (2013). Optimization and evaluation of a multiplex PCR for simultaneous detection of prominent foodborne pathogens of *Enterobacteriaceae*. *Annals of Microbiology*, 63(4), 1591-1599.
- Brise, S., Fevre, C., Passet, V., Issenhuth-Jeanjean, S., Tournebize, R., Diancourt, L., & Grimont, P. (2009). Virulent clones of *Klebsiella pneumoniae*: identification and evolutionary scenario based on genomic and phenotypic characterization. *PLoS One*, 4(3), 4982.
- Eibach, D., Dekker, D., Boahen, K. G., Akenten, C. W., Sarpong, N., Campos, C. B., & May, J. (2018). Extended-spectrum beta-lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* in local and imported poultry meat in Ghana. *Veterinary microbiology*, 217, 7-12.
- Fielding, B. C., Mnabisa, A., Gouws, P. A., & Morris, T. (2012). Antimicrobial-resistant *Klebsiella* species isolated from free-range chicken samples in an informal settlement. *Archives of Medical Science: Archives of Medical Science*, 8 (1), 39-42.
- Gundogan, N., Citak, S., & Yalcin, E. (2011). Virulence Properties of Extended Spectrum  $\beta$ -Lactamase-Producing *Klebsiella* Species in Meat Samples. *Journal of food protection*, 74(4), 559-564.
- Guo, Y., Zhou, H., Qin, L., Pang, Z., Qin, T., Ren, H., & Zhou, J. (2016). Frequency, antimicrobial resistance and genetic diversity of *Klebsiella pneumoniae* in food samples. *PLoS One*, 11(4), 0153561.
- Hayati, M., Indrawati, A., Mayasari, N. L. P. I., Istiyaningsih, I., & Atikah, N. (2019). Molecular detection of extended-spectrum  $\beta$ -lactamase-producing *Klebsiella pneumoniae* isolates of chicken origin from East Java, Indonesia. *Veterinary World*, 12(4), 578-583.
- Hermana, N. S. P., Nilasari, Z., Yuniarti, E., Riyanto, A., Indrawati, A., & Pasaribu,

- F. H. (2020, January). Antibiotic resistance profile of *Staphylococcus aureus*, *Streptococcus* spp. and *Klebsiella* spp. isolated from chicken farm in Bogor, Sukabumi, and Cianjur, West Java. In *Journal of Physics: Conference Series*. 1430, 012021.
- Janda, J. M., & Abbott, S. L. (2007). 16S rRNA gene sequencing for bacterial identification in the diagnostic laboratory: pluses, perils, and pitfalls. *Journal of Clinical Microbiology*, 45(9), 2761-2764.
- Kurupati, P., Chow, C., Kumarasinghe, G., & Poh, C. L. (2004). Rapid detection of *Klebsiella pneumoniae* from blood culture bottles by real-time PCR. *Journal of clinical microbiology*, 42(3), 1337-1340.
- Mahanti, A., Ghosh, P., Samanta, I., Joardar, S. N., Bandyopadhyay, S., Bhattacharyya, D., & Dutta, T. K. (2018). Prevalence of CTX-M-Producing *Klebsiella* spp. in broiler, kuroiler, and indigenous poultry in West Bengal State, India. *Microbial Drug Resistance*, 24(3), 299-306.
- National Action Plan for Egg & Poultry, (2017). National Action Plan for Egg & Poultry-2022 for Doubling Farmers' Income by 2022. Retrieved from [http://www.dahd.nic.in/sites/default/files/Seeking%20Comments%20on%20National%20Action%20Plan-%20Poultry-%202022%20by%2012-12-2017\\_0\\_0.pdf](http://www.dahd.nic.in/sites/default/files/Seeking%20Comments%20on%20National%20Action%20Plan-%20Poultry-%202022%20by%2012-12-2017_0_0.pdf).
- Rajaa, M., Milyani, Khadija, Al-Aali, Y. (2011). Enterobacteria Carried by *Musca domestica* in the Western Region of Saudi Arabia. *J. Environ. Applicat. Sci.* 6(2), 290–298.
- Razmyar, J., & Zamani, A. H. (2016). An outbreak of yolk sac infection and dead-in-shell mortality in common canary (*Serinus canaria*) caused by *Klebsiella pneumoniae*. *Iranian journal of veterinary research*, 17(2), 141-143.
- Sękowska, A. (2017). *Raoultella* spp. clinical significance, infections and susceptibility to antibiotics. *Folia Microbiologica*, 62(3), 221-227.
- Veeraselvam, M., Senthilkumar, N. R., Vairamuthu, S., & Ramakrishnan, V. (2019). Isolation and identification of bacterial agents causing respiratory infection in native chicken. *Journal of Entomology and Zoology Studies*, 7(4), 162-167.
- Von Tippelskirch, P., Gölz, G., Projahn, M., Daehre, K., Friese, A., Roesler, U., & Orquera, S. (2018). Prevalence and quantitative analysis of ESBL and AmpC beta-lactamase producing *Enterobacteriaceae* in broiler chicken during slaughter in Germany. *International journal of food microbiology*, 281, 82-89.
- Wu, H., Wang, M., Liu, Y., Wang, X., Wang, Y., Lu, J., & Xu, H. (2016). Characterization of antimicrobial resistance in *Klebsiella* species isolated from chicken broilers. *International Journal of Food Microbiology*, 232(9), 95-102.
- Younis, G., Awad, A., El-Gamal, A., & Hosni, R. (2016). Virulence properties and antimicrobial susceptibility profiles of *Klebsiella* species recovered from clinically diseased broiler chicken. *Advances in Animal and Veterinary Sciences*, 4(10), 536-542.